

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
3 April 2008 (03.04.2008)

PCT

(10) International Publication Number  
**WO 2008/038062 A1**

(51) International Patent Classification:

*C12N 1/20* (2006.01) *C12Q 1/68* (2006.01)  
*C12N 15/09* (2006.01)

**Jacques** [FR/FR]; 8, Chemin de Versailles, F-34980 Montferrier-sur-Lez (FR). **MARTINEZ, Dominique** [FR/FR]; 5, Route de Quissac, F-30610 Sauve (FR).

(21) International Application Number:

PCT/IB2006/003870

(74) Agents: **VIALLE-PRESLES, Marie-José** et al.; Cabinet Ores, 36, rue de Saint-Petersbourg, F-75008 Paris (FR).

(22) International Filing Date:

25 September 2006 (25.09.2006)

(25) Filing Language:

English

(26) Publication Language:

English

(71) Applicants (for all designated States except US): **CENTRE DE COOPERATION INTERNATIONALE EN RECHERCHE AGRONOMIQUE POUR LE DEVELOPPEMENT (CIRAD)** [FR/FR]; Etablissement Public à caractère Industriel et Commercial, 42, rue Scheffer, F-75016 Paris (FR). **CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE** [FR/FR]; Etablissement Public, 3, rue Michel Ange, F-75016 Paris (FR).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **FRUTOS, Roger** [FR/FR]; 1, rue des Alicantes, F-34680 St. Georges d'Orques (FR). **VACHIERY, Nathalie** [FR/FR]; Chemin de Ravine Chaude, F-97129 Lamentin (FR). **LEFRANCOIS, Thierry** [FR/FR]; 26, lot les Belles Vues de Convenance, F-97122 Baie Mahault (FR). **FERRAZ, Conception** [FR/FR]; Le Clos des Elbes, 114, rue des Eucalyptus, F-34090 Montpellier (FR). **DEMAILLE,**

Published:

- with international search report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

(54) Title: TARGET GENES FOR STRAIN-SPECIFIC DIAGNOSTIC OF EHRLICHIA RUMINANTIIUM AND USE THEREOF

(57) Abstract: The invention provides a combination of target genes that are useful as genetic markers for the strain-specific detection of *Ehrlichia ruminantium*. The invention also provides diagnostic methods using said combination of markers.



WO 2008/038062 A1

## TARGET GENES FOR STRAIN-SPECIFIC DIAGNOSTIC OF *EHRlichia RUMINANTIUM* AND USE THEREOF

*Rickettsiae* are intracellular pathogenic bacteria responsible for various diseases on Humans and animals. *Rickettsiae* are transmitted by arthropods, most frequently ticks, lice and mites, and cause major illnesses such as epidemic typhus or Rocky Mountain spotted fever. The genus *Ehrlichia* comprises other species pathogenic for humans and mammals such as *E. chaffeensis*, responsible for Human monocytic ehrlichiosis, *E. canis*, the causing agent of canine monocytic ehrlichiosis.

Another species, *Ehrlichia ruminantium*, formerly known as *Cowdria ruminantium*, is the causing agent of heartwater or cowdriosis, an economically important disease of domestic ruminants. Heartwater can cause up to 80 % mortality in susceptible animals. *E. ruminantium* is transmitted by *Amblyomma* ticks and is present in Sub-Saharan Africa and surrounding islands, including Madagascar. Heartwater is also present in several Caribbean islands and is threatening the American mainland.

Vaccination against heartwater has long been based on "infection and treatment". Naïve animals are inoculated with blood containing virulent organisms, a procedure which carries a high risk of uncontrolled clinical reactions and the inadvertent spread of undesirable parasites and viruses. A first generation cowdriosis inactivated vaccine based on cell-cultured derived elementary bodies was developed. Although representing a considerable improvement and the first heartwater vaccine acceptable for widespread use, the level of protection conferred is still not fully satisfactory. Indeed, all animals develop a clinical reaction at challenge despite vaccination. Furthermore, livestock also faces challenge by genetically and antigenically diverse strains.

Diversity of *E. ruminantium* is a key problem which has been recognized for a long time, but insufficient information is available for optimum vaccine formulation and specific diagnostic. Serological diagnostic tests of heartwater using crude antigens from whole bacteria detect false positive reactions due to common antigenic determinants

The diversity of *E. ruminantium* was demonstrated at the antigenic level by cross-immunisation studies. Variable antigens were identified by ELISA and immunoblot using cross-absorbed immune sera.

Genetic diversity was later demonstrated when sequencing the Map 1 gene which showed a high degree of sequence heterogeneity concentrated in three hypervariable regions. Genomic polymorphism was also detected using RAPD and RFLP markers. This DNA polymorphism was shown to correlate with antigenic polymorphism.

ELISA-based and serological diagnostics have been developed using the Map 1 and the GroEL (WO 9914233) antigens. Other peptides for serological diagnostic have been described (US 2002004051, US 20020132789, WO 02/066652). Although they have dramatically improved specificity, they still display cross reaction with *E. canis* and *E.*

*chaffeensis*. The *map1* gene initially considered as a good marker for geographic diversity, was recently shown not to be geographically constrained. Furthermore, the life span of anti-Map 1 antibodies is rather short.

PCR-based diagnostic methods represent methods of choice for the sensitive and specific detection of *Ehrlichia* in clinically reactive or asymptomatic carrier ruminants, as well as in vectors. However, in the field, hosts and vectors can be co-infested by several parasites and the diversity of pathogen species is further complicated by the existence of extensive intra-species diversity. Thus, it is important to provide means and diagnostic tools allowing not only to identify *E. ruminantium* but also to differentiate between different strains.

Sequences allowing differential diagnostic of *E. ruminantium* strain Gardel and *E. ruminantium* strain Welgevonden have been previously described by the inventors. They have shown, through complete genome sequencing and comparative genomic analysis that several genes were only found in either strain Gardel or strain Welgevonden, without counterpart in the other strain, and that several other genes, while being present in both strains differed between them by one or several mutations, such as large insertions and/or deletions that result in a frameshift and/or in a truncated version of the original gene. These genes were therefore primary targets to develop specific, multitarget diagnostic methods to differentiate between these two strains (WO 2006/045338; Frutos et al., Journal of Bacteriology.188: 2533-2542, 2006).

The inventors have now found that the use of a particular combination of some of the target genes described in WO 2006/045338 allowed not only to discriminate between strains Gardel and Welgevonden, but also in a more general way, to detect specifically *E. ruminantium* and to discriminate between a broad range of strains of *E. ruminantium* other than Gardel and Welgevonden including strains for which no genomic sequence data are available.

An object of the invention is thus the use of the following set of genes:

Erum1, defined by the sequence SEQ ID NO: 6

Erum2, defined by the sequence SEQ ID NO: 3

Erum3, defined by the sequence SEQ ID NO: 1

Erum4, defined by the sequence SEQ ID NO: 4

Erum5, defined by the sequence SEQ ID NO: 2

Erum6, defined by the sequence SEQ ID NO: 5

Erum7, defined by the sequence SEQ ID NO: 13

Erum8, defined by the sequence SEQ ID NO: 15

Erum9, defined by the sequence SEQ ID NO: 14

Erum10, defined by the sequence SEQ ID NO: 8,

as targets for the strain-specific detection of *Ehrlichia ruminantium*.

The reference sequences used herein to define the target genes Erum 1-5 and Erum 7-9 are those identified in the Gardel strain; the reference sequences used herein to define the target genes Erum6 and Erum10 are those identified in the Welgevonden strain.

5 However, it is to be understood that each of these genes actually exists under different allelic forms, depending on the strain of *Ehrlichia ruminantium*. The allelic forms that will be considered herein, having in view strain-specific detection, are in particular those resulting from large insertions and/or deletions that lead to a frameshift or to a truncated version of the original gene.

10 The invention thus provides a method for the strain-specific detection of *Ehrlichia ruminantium* wherein said method comprises determining, for each of the genes Erum 1 to Erum10 defined above, whether said gene is present in the bacteria to be tested, and under which allelic form.

15 Advantageously, the method of the invention is carried out by performing PCR amplification of all the target genes Erum 1 to Erum10, and checking, for each of these genes, the presence of one or more amplification product(s), and the size of said amplification product(s).

Within the target genes Erum 1 to Erum10, preferred target regions are as follows:

20 For Erum 1, the target region can consist of the whole sequence SEQ ID NO: 6, or of a portion thereof; in particular the target region can be defined within the portion spanning from nucleotide 1 to nucleotide 173 of SEQ ID NO: 6.

For Erum 2, the target region can consist of the whole sequence SEQ ID NO: 3, or of a portion thereof; in particular the target region can be defined within the portion spanning from nucleotide 1 to nucleotide 218 of SEQ ID NO: 3.

25 For Erum 3, the target region can consist of the whole sequence SEQ ID NO: 1, or of a portion thereof; in particular the target region can be defined within the portion spanning from nucleotide 1 to nucleotide 509 of SEQ ID NO: 1.

30 For Erum 4, the target region can consist of the whole sequence SEQ ID NO: 4, or of a portion thereof; in particular the target region can be defined within the portion spanning from nucleotide 56 to nucleotide 698 of SEQ ID NO: 4.

For Erum 5, the target region can consist of the whole sequence SEQ ID NO: 2, or of a portion thereof; in particular the target region can be defined within the portion spanning from nucleotide 1 to nucleotide 239 of SEQ ID NO: 2.

35 For Erum 6, the target region can consist of the whole sequence SEQ ID NO: 5, or of a portion thereof; in particular the target region can be defined within the portion spanning from nucleotide 3 to nucleotide 130 of SEQ ID NO: 5.

For Erum 7, a preferred target region is located within the portion spanning from nucleotide 1 to nucleotide 1981 of SEQ ID NO: 13; another preferred target region is

located within the portion spanning from nucleotide 2378 to nucleotide 3252 of SEQ ID NO: 13.

For Erum 8, a preferred target region is located within the portion spanning from nucleotide 1 to nucleotide 926 of SEQ ID NO: 15; another preferred target region is  
5 located within the portion spanning from nucleotide 1816 to nucleotide 3570 of SEQ ID NO: 15.

For Erum 9, a preferred target region is located within the portion spanning from nucleotide 1 to nucleotide 1307 of SEQ ID NO: 14; another preferred target region is located within the portion spanning from nucleotide 151 to nucleotide 1836 of  
10 SEQ ID NO: 14.

For Erum 10, a preferred target region is located within the portion spanning from nucleotide 1 to nucleotide 598 of SEQ ID NO: 8; another preferred target region is located within the portion spanning from nucleotide 792 to nucleotide 3522 of SEQ ID NO: 8; still another target region is located within the portion spanning from nucleotide 599 to  
15 nucleotide 791 of SEQ ID NO: 8.

Various techniques for detection of target nucleic acid sequences based on PCR amplification are available in the art.

These methods include in particular combined PCR analysis, i.e. simultaneous gel visualization of ten individual PCR reactions, each one targeting only one of  
20 the genes Erum1 to Erum10 defined above. The ten target genes can also be analysed by multiplex PCR, by a single PCR reaction involving simultaneous amplification of all the genes using a mixture of primers and visualization of the pattern on electrophoresis gel, or by a combination of multiplex PCR reactions, each one concerning a subset of the target genes listed above.

Non-limitative examples of PCR primers allowing to carry out the method of the invention are given in Table 2 below. Other suitable PCR primers can easily be designed by one of skill in the art, on the basis of the information provided by the present invention. By way of non-limitative example of oligonucleotide design software suitable for obtaining PCR primers of the invention, one can mention the software Vector NTI Advance  
30 9.0 (Invitrogene).

The invention also comprises diagnostic kits for discriminating between strains of *E. ruminantium* wherein said kits comprise PCR primers for all the target genes Erum 1 to Erum10.

The method of the invention is useful in particular to discriminate between  
35 strains of *E. ruminantium* other than strain Gardel and strain Welgevonden. It is also useful to discriminate between strain Gardel and strains of *E. ruminantium* other than strain Welgevonden, or conversely, between strain Welgevonden and strains of *E. ruminantium*

other than strain Gardel. Furthermore, it also allows for discriminating between a virulent strain of *E. ruminantium* and its attenuated counterpart.

The method of the invention can be performed either on whole bacteria previously lysed, or on nucleic acid (genomic DNA, cDNA or mRNA) isolated from said bacteria. It is suitable for use at various stages of the life cycle of *E. ruminantium*, more specifically but not limited to the domestic-ruminants infectious stage, vector-interaction stage or reservoir animals-interaction stage. Preferred utilisations of the method of the invention include the detection of *Ehrlichia ruminantium* in a given territory, the strain specific identification of *Ehrlichia ruminantium* in a given territory, the discrimination between strains of *Ehrlichia ruminantium* in a given territory or between different geographical regions, the analysis of strain movements within a region or between geographically distinct regions, the differential presence of strains of *Ehrlichia ruminantium* according to vector species and/or populations or the early detection and risk assessment in regions where potential vectors are present but where the disease has not been recorded yet.

Specifically exemplified herein is the identification of *E. ruminantium* strains based on the specific amplification patterns of the ten target genes defined above,

#### EXAMPLE 1. GENERAL FEATURES AND SEQUENCE REFERENCE

For each strain, purified DNA was broken by sonication to generate fragments of differing sizes. After filling up the ends with Klenow polymerase, DNA fragments ranging from 0.5 kb to 4 kb were separated in a 0.8% agarose gel and collected after gelase (Epicentre) digestion of a cut agarose band. Blunt-end DNA fragments were inserted into pBluescript II KS (Stratagene) digested with EcoRV and dephosphorylated. Ligation was performed with the Fast-Link DNA Ligation kit (Epicentre) and competent DH10B *E. coli* were transformed prior to colony isolation on LB-agar+ Ampicillin + Xgal +IPTG. About 15000 clones were isolated for each strain of *E. ruminantium*. Plasmidic DNA from recombinant *E. coli* strains was extracted according to the alkaline lysis method and inserts were sequenced on both strands using universal forward and reverse M13 primers and the ET DYEnamic terminator kit (Amersham). Sequences were obtained with ABI 373 et ABI 377 automated sequencers (Applied Biosystems). Data were analysed and contigs were assembled using Phred-Phrap and Consed software packages (<http://www.genome.washington.edu>). Gaps were filled in through primer-directed sequencing using custom made primers. A total of about 20000 raw sequence runs were generated and analysed for each *E. ruminantium* strain to generate a full length consensus sequence with a coverage of 6x to 7x.

*E. ruminantium* strain Gardel and *E. ruminantium* strain Welgevonden are virulent pathogenic strains causing heartwater in Guadeloupe Island (French West Indies) and South Africa, respectively. The genome of *E. ruminantium* strains Gardel and Welgevonden is arranged as a circular chromosome of 1499920 bp and 1512977 bp, respectively. The

respective G+C contents for the strains Gardel and Welgevonden is 27.51 % and 27.48 %. The genome of *E. ruminantium* strain Gardel comprises 948 coding sequences of an average size of 1018 bp which represent a total coding surface of 63 % of the whole genome. The genome of *E. ruminantium* strain Welgevonden bears 957 genes of the same average size of 1018 bp. The genome surface of this strain devoted to coding sequences is 62 %. Both genomes comprise 36 transfer RNAs (tRNA) and 3 ribosomal RNAs (rRNA).

**EXAMPLE 2. IDENTIFICATION OF TARGET GENES FOR STRAIN SPECIFIC DIFFERENTIAL DIAGNOSTIC IN THE GARDEL AND WELGEVONDEN STRAINS OF *E. RUMINANTIUM***

The differential analysis of the whole genomes of *E. ruminantium* strains Gardel and Welgevonden showed the presence of coding sequences which are present in only one of the strains and not in the other. Some of the CDS which are unique to *E. ruminantium* strain Gardel and found only in the genome of this strain are presented in Table 1 (Seq ID NO 1 to Seq ID NO 5). One of the CDS which is unique to *E. ruminantium* strain Welgevonden and found only in the genome of this strain is presented in Table 1 (Seq ID NO 6). Since these sequences are unique to one or the other strain, they clearly represent targets for the differential detection of *E. ruminantium* strain Gardel versus *E. ruminantium* strain Welgevonden.

The differential analysis of the whole genomes of *E. ruminantium* strains Gardel and Welgevonden also showed the presence of coding sequences which are affected by one or several mutations in one of the two strains and for which a non-mutated, functionally active and normal allele is present in the genome of the other strain. Mutations yielded a stop codon which may result in shorter but still predicted CDS depending upon the size of the remaining fragments. Truncated genes resulting in a single CDS are denominated partial CDS, whereas those resulting in two or more predicted CDS are described as fragmented CDS. These coding sequences are presented in Table 1. One Such CDS in the genome of *E. ruminantium* strain Gardel which is affected by mutations and differs from its native counterpart in *E. ruminantium* strain Welgevonden is presented in Table 1 (SEQ ID NO 7). This is a truncated version of the native gene in *E. ruminantium* strain Welgevonden (Table 1, SEQ ID NO 8). The genome of *E. ruminantium* strain Welgevonden also bears mutated genes, with respect to their allelic variant counterparts in the genome of *E. ruminantium* strain Gardel. Three of these CDS which are affected by mutations generating a truncated version of the genes are presented in Table 1 (SEQ ID NO 9 to SEQ ID NO 12). The native full length allele of these CDS present in the genome of *E. ruminantium* strain Gardel are shown in Table 1 (SEQ ID NO 13 to SEQ ID NO 15). One series of CDS in *E. ruminantium* strain Welgevonden (SEQ ID NO 11 and SEQ ID NO 12), whose native full length alleles are found in the genome of *E. ruminantium* strain Gardel (Table 1, SEQ ID NO 15) was affected by mutations generating a frameshift.

Table 1. Target genes for strain-specific differential diagnostic of *E. ruminantium*

Target gene	Gene in Gardel	Status	Gene in Welgevonden	Status
Erum 3	ERGA_CDS_05600 (SEQ ID No 1)	Unique gene	None	Sequence absent (full deletion)
Erum 5	ERGA_CDS_07600 (SEQ ID No 2)	Unique gene	None	Sequence absent (full deletion)
Erum 2	ERGA_CDS_04990 (SEQ ID No 3)	Unique gene	None	Partial deletion
Erum 4	ERGA_CDS_05610 (SEQ ID No 4)	Unique gene	None	Extensive mutations
Erum 6	None	Partial deletion	ERWE_CDS_08340 (SEQ ID No 5)	Unique gene
Erum 1	ERGA_CDS_04350 (SEQ ID No 6)	Unique gene	None	Extensive mutations
Erum 10	ERGA_CDS_07340 (SEQ ID No 7)	Partial deletion	ERWE_CDS_07420 (SEQ ID No 8)	Full length gene
Erum 7	ERGA_CDS_01350 (SEQ ID No 13)	Full length gene	ERWE_CDS_01390 (SEQ ID No 9)	Partial deletion
Erum 9	ERGA_CDS_05750 (SEQ ID No 14)	Full length gene	ERWE_CDS_05840 (SEQ ID No 10)	Partial deletion
Erum 8	ERGA_CDS_04510 (SEQ ID No 15)	Full length gene	ERWE_CDS_04590 (SEQ ID No 11) ERWE_CDS_04600 (SEQ ID No 12)	Frameshift (partial deletion)

**EXAMPLE 3. DIFFERENTIAL DETECTION OF STRAIN GARDEL AND STRAIN WELGEVONDEN OF *E. RUMINANTIUM* BASED ON PCR AMPLIFICATION PATTERNS OF THE TARGET GENES**

5 Differential PCR identification of strains Gardel and Welgevonden of *E. ruminantium* was achieved using primers described in Table 2.



Table 2

Target gene	Primer name	SEQ ID	Orientation	Size (mer)	CDS
Erum 1	P-Erum 1-A	SEQ ID #16	Sense	21	ERGA_CDS_04350
Erum 1	P- Erum 1-B	SEQ ID #17	Antisense	21	ERGA_CDS_04350
Erum 2	P- Erum 2-A	SEQ ID #18	Sense	25	ERGA_CDS_4990
Erum 2	P- Erum 2-B	SEQ ID #19	Antisense	23	ERGA_CDS_4990
Erum 3	P- Erum 3-A	SEQ ID #20	Sense	20	ERGA_CDS_05600
Erum 3	P- Erum 3-B	SEQ ID #21	Antisense	20	ERGA_CDS_05600
Erum 4	P- Erum 4-A	SEQ ID #22	Sense	19	ERGA_CDS_05610
Erum 4	P- Erum 4-B	SEQ ID #23	Antisense	22	ERGA_CDS_05610
Erum 5	P- Erum 5-A	SEQ ID #24	Sense	23	ERGA_CDS_07600
Erum 5	P- Erum 5-B	SEQ ID #25	Antisense	19	ERGA_CDS_07600
Erum 6	P- Erum 6-A	SEQ ID #26	Sense	26	ERWE_CDS_08340
Erum 6	P- Erum 6-B	SEQ ID #27	Antisense	23	ERWE_CDS_08340
Erum 7	P- Erum 7-A	SEQ ID #28	Sense	25	ERGA_CDS_01350 ERWE_CDS_01390
Erum 7	P- Erum 7-B	SEQ ID #29	Antisense	25	ERGA_CDS_01350 ERWE_CDS_01390 ERGA_CDS_04510
Erum 8	P- Erum 8-A	SEQ ID #30	Sense	25	ERWE_CDS_04590 ERWE_CDS_04600 ERGA_CDS_04510
Erum 8	P- Erum 8-B	SEQ ID #31	Antisense	25	ERWE_CDS_04590 ERWE_CDS_04600 ERGA_CDS_05750
Erum 9	P- Erum 9-A	SEQ ID #32	Sense	25	ERWE_CDS_05840 ERGA_CDS_05750
Erum 9	P- Erum 9-B	SEQ ID #33	Antisense	25	ERWE_CDS_05840 ERGA_CDS_07340
Erum 10	P- Erum 10-A	SEQ ID #34	Sense	25	ERWE_CDS_07420
Erum 10	P- Erum 10-B	SEQ ID #35	Antisense	25	ERGA_CDS_07340 ERWE_CDS_07420

DNA is extracted from elementary bodies of *E. ruminantium*, as described by Perez *et al.* (1997). *E. ruminantium* strains are grown in BUEC cells as described above. Elementary bodies are purified from the culture supernatant by differential centrifugation and resuspended in 350  $\mu$ l of PBS to which is added 150  $\mu$ l of buffer containing 25 mM Tris-HCl (pH 8.0), 10 mM MgCl<sub>2</sub> and 125  $\mu$ g of DNase in order to remove contaminating host cell DNA. After incubation for 90 min. at 37°C, the reaction is stopped by addition of 25 mM EDTA. Elementary bodies are washed three times in water and lysed by overnight incubation at 55°C in a solution of 100 mM Tris-HCl (pH 8.0), 150 mM NaCl, 25 mM EDTA, 1.5% SDS and 250  $\mu$ g/ml of proteinase K. Bacterial DNA is extracted with phenol-choloroform,

precipitated with cold ethanol and resuspended in sterile distilled water. Contamination with cell DNA is evaluated by slot blot hybridization using labeled bovine DNA as a probe and dilutions of bovine DNA (12.5 ng and 25 ng) as positive controls.

5 PCR amplification of amplicons is performed by mixing 250 ng of *E. ruminantium* DNA, 2.5 U of Taq DNA polymerase, 200 nM of each dNTP, 1 µM of each, sense and antisense, primer and 3 mM MgCl<sub>2</sub> in a final volume of 50 µl. Amplification is done under the following conditions: 5 min denaturation at 94°C, followed by 30 cycles of amplification with a 1-min denaturation, 45 sec of annealing at 45°C and 2 min extension at 72°C. An extra extension step of 10 min at 72°C is added after completion of the 30 cycles. PCR products, i.e. amplicons, are analysed by 1% agarose gel electrophoresis in Tris-borate-EDTA buffer.

The results are summarized in Table 3, Figure 1, and Figure 2.

#### Legend of Figure 1:

1: Molecular weight marker (100-bp ladder); 2: *Erga* with primers P-Erum 1-A + P-Erum 1-B; 3: *Erwe* with primers P-Erum 1-A + P-Erum 1-B; 4: Control sample with primers P-Erum 1-A + P-Erum 1-B; 5: *Erga* with primers P-Erum 2-A + P-Erum 2-B; 6: *Erwe* with primers P-Erum 2-A + P-Erum 2-B; 7: Control sample with primers P-Erum 2-A + P-Erum 2-B; 8: *Erga* with primers P-Erum 3-A + P-Erum 3-B; 9: *Erwe* with primers P-Erum 3-A + P-Erum 3-B; 10: Control sample with primers P-Erum 3-A + P-Erum 3-B; 11: *Erga* with primers P-Erum 4-A + P-Erum 4-B; 12: *Erwe* with primers P-Erum 4-A + P-Erum 4-B; 13: Control sample with primers P-Erum 4-A + P-Erum 4-B; 14: *Erga* with primers P-Erum 5-A + P-Erum 5-B; 15: *Erwe* with primers P-Erum 5-A + P-Erum 5-B; 16: Control sample with primers P-Erum 5-A + P-Erum 5-B; 17: *Erga* with primers P-Erum 6-A + P-Erum 6-B; 18: *Erwe* with primers P-Erum 6-A + P-Erum 6-B; 19: Control sample with primers P-Erum 6-A + P-Erum 6-B; 20: Molecular weight marker (1 *Eco*RI-*Hind*III).

#### Legend of Figure 2:

1: Molecular weight marker (100-bp ladder); 2: *Erga* with primers P-Erum 7-A + P-Erum 7-B; 3: *Erwe* with primers P-Erum 7-A + P-Erum 7-B; 4: Control sample with primers P-Erum 7-A + P-Erum 7-B; 5: *Erga* with primers P-Erum 9-A + P-Erum 9-B; 6: *Erwe* with primers P-Erum 9-A + P-Erum 9-B; 7: Control sample with primers P-Erum 9-A + P-Erum 9-B; 8: *Erga* with primers P-Erum 10-A + P-Erum 10-B; 9: *Erwe* with primers P-Erum 10-A + P-Erum 10-B; 10: Control sample with primers P-Erum 10-A + P-Erum 10-B; 11: Molecular weight marker (1 *Eco*RI-*Hind*III); 12: Molecular weight marker (100-bp ladder); 13: *Erga* with primers P-Erum 8-A + P-Erum 8-B; 14: *Erwe* with primers P-Erum 8-A + P-Erum 8-B; 15: Control sample with primers P-Erum 8-A + P-Erum 8-B

As shown in Table 3, Fig.1 and Fig. 2, all the pairs of primers described in Table 3 allowed for differential identification and discrimination of strains Gardel and Welgevonden. The PCR reactions yielded the results expected from *in silico* prediction of

amplicons (Table 3). The pairs P-Erum 1-A + P-Erum 1-B, P-Erum 2-A + P-Erum 2-B, P-Erum 3-A + P-Erum 3-B, P-Erum 4-A + P-Erum 4-B, P-Erum 5-A + P-Erum 5-B, which target unique genes present only in strain Gardel generated the expected unique amplicons of 172 bp, 217 bp, 508 bp, 642 bp and 238 bp, respectively, on strain Gardel while generating no bands on strain Welgevonden (Table 3, Fig. 1). The pair of primers P-Erum 6-A + P-Erum 6-B which target a unique gene only present in strain Welgevonden yielded, as expected, a single amplicon of 127 bp while no PCR product was obtained on strain Gardel (Table 3, Fig. 1). The pairs P-Erum 7-A + P-Erum 7-B, P-Erum 8-A + P-Erum 8-B, P-Erum 9-A + P-Erum 7-B and P-Erum 10-A + P-Erum 10-B targeting the truncated genes yielded PCR products of the respective expected size of 2791 bp, 552 bp+ 1071 bp, 1361 bp and 1095 bp on strain Gardel and 2395 bp, 492 bp, 1178 bp and 1691 bp on strain Welgevonden, respectively (Table 3, Fig. 2). An additional band of 480 bp is observed on strain Gardel with the pair P-Erum 8-A + P-Erum 8-B. This additional band is most likely due to a single low specificity response occurring in *Erga*.

The primer pairs P-1350-A + P-1350-B, P-4510-A + P-4510-B, P-5750-A + P-5750-B and P-7420-A + P-7420-B yielded PCR products of the respective expected size of 2791, 552+1071, 1361 and 1095 on strain Gardel and 2395, 492, 1178 and 1691 on strain Welgevonden, respectively (Table 3, Fig. 1 and Fig. 2). An additional band of 480 bp is observed on strain Gardel with the pair P-4510-A + P-4510-B. This additional band is most likely due to a single low specificity response occurring in strain Gardel.

Table 3. Strain-specific differential PCR screening of *E. ruminantium* strain Gardel and strain Welgevonden

Primer combination	Strain			
	Gardel		Welgevonden	
	Expected	Observed	Expected	Observed
P-Erum 1-A + P-Erum 1-B	172	172	None	None
P-Erum 2-A + P-Erum 2-B	217	217	None	None
P-Erum 3-A + P-Erum 3-B	508	508	None	None
P-Erum 4-A + P-Erum 4-B	642	642	None	None
P-Erum 5-A + P-Erum 5-B	238	238	None	None
P-Erum 6-A + P-Erum 6-B	None	None	127	127
P-Erum 7-A + P-Erum 7-B	2791	2791	2395	2395
P-Erum 8-A + P-Erum 8-B	552+1071	552+1071+ 480	492	492
P-Erum 9-A + P-Erum 9-B	1361	1361	1178	1178
P-Erum 10-A + P-Erum 10-B	1095	1095+300	1691	1691

#### EXAMPLE 4: DIFFERENTIAL STRAIN-SPECIFIC PCR DETECTION AND IDENTIFICATION OF STRAINS OF *E. RUMINANTIUM* DIFFERENT THAN STRAIN GARDEL AND STRAIN WELGEVONDEN

The use primers listed in Table 2 were used for the specific identification and discrimination of *E. ruminantium* strains other than strain Gardel and strain Welgevonden. The strains others than Gardel and Welgevonden presented in this example are strains Umpala (Mozambique), Senegal (Senegal), Bankouma (Burkina Faso) , Bekuy (Burkina Faso), Lamba (Burkina Faso), Banan 1 (Burkina Faso) and Banan 2 (Burkina Faso).

These strains are presented here to illustrate samples from different parts of Sub-Saharan Africa and the Caribbean.

DNA is extracted from elementary bodies of *E. ruminantium* and PCR amplification performed as described in Example 3.

5 The results are shown in Table 4, Fig. 3 and Fig. 4:

Legend of Figure 3:

- A. PCR detection with primers P-Erum 1-A + P-Erum 1-B
- B. PCR detection with primers P-Erum 3-A + P-Erum 3-B
- C. PCR detection with primers P-Erum 2-A + P-Erum 2-B
- 10 D. PCR detection with primers P-Erum 4-A + P-Erum 4-B
- E. PCR detection with primers P-Erum 6-A + P-Erum 6-B
- F. PCR detection with primers P-Erum 5-A + P-Erum 5-B

MW1: Molecular weight marker (100pb DNA ladder); MW2: Molecular weight marker (1  
HindIII/EcoRI); 1: Strain Senegal attenuated (Satt); 2: Strain Gardel CTVM; 3: Strain  
15 Bankouma; 4: Strain Bekuy; 5: Strain Lamba, 6: Strain Banan 1; 7: Strain Banan 2; NC:  
Negative control; G. Strain Gardel; W: Strain Welgevonden.

Legend of Figure 4:

- A. PCR detection with primers P-Erum 7-A + P-Erum 7-B
- B. PCR detection with primers P-Erum 8-A + P-Erum 8-B
- 20 C. PCR detection with primers P-Erum 9-A + P-Erum 9-B
- D. PCR detection with primers P-Erum 10-A + P-Erum 10-B

MW1: Molecular weight marker (100pb DNA ladder); MW2: Molecular weight marker (1  
HindIII/EcoRI); 1: Strain Bankouma; 2: Strain Bekuy; 3: Strain Lamba; 4: Strain Banan 1; 5:  
Strain Banan 2; 6: Strain Gardel attenuated (Gatt); 7: Strain Gardel CTVM; 8: Strain  
25 Senegal; 9: Strain Senegal attenuated (Satt); NC: Negative control; G. Strain Gardel; W:  
Strain Welgevonden.

As shown in Table 4, Fig. 3 and Fig. 4, the combined use of all the pairs of  
primers described in Table 2 allowed for differential identification and discrimination of  
strains other than strains Gardel and Welgevonden. The PCR reactions results are summarized  
30 in Table 4. The pairs P-Erum 1-A + P-Erum 1-B, P-Erum 2-A + P-Erum 2-B, P-Erum 3-A +  
P-Erum 3-B, P-Erum 4-A + P-Erum 4-B, P-Erum 5-A + P-Erum 5-B which target unique  
genes present only in strain Gardel and the pair P-Erum 6-A + P-Erum 6-B which targets a  
unique gene only present in strain Welgevonden all yielded differing patterns of PCR  
products depending on the strain (Table 4, Fig 3). Differing patterns depending upon the  
35 strain were also observed using the pairs P-Erum 7-A + P-Erum 7-B, P-Erum 8-A + P-Erum  
8-B, P-Erum 9-A + P-Erum 7-B and P-Erum 10-A + P-Erum 10-B which target the truncated  
genes (Table 4, Fig. 3).

It is however the overall analysis of all the PCR patterns yielded by all the pairs of primers described in Table 2 which provides a strain specific diagnostic. The strains Bekuy and Lamba which were isolated in Burkina Faso from the nearby villages of Bekuy and Lamba, respectively, are most likely to be two isolates of the same strain. Furthermore, these strains display the same map-1 genotype determined by PCR amplification and sequencing of the map-1 gene. All the other strains display differing map-1 genotypes. This further indicates that strains Bekuy and Lamba are two isolate of the same strain. The identical overall pattern obtained for these two strains with all the pairs of primers described in Table 2 also further demonstrate the strain-specificity of the subject of the invention and its ability to identify different strains and separate isolates of the same strain.

Table 4: Strain-specific differential PCR screening of *E. ruminantium*

Primer combination	Strain								
	Gardel	Welgevonden	Umpala	Senegal	Bankouma	Bekuy	Lamba	Banan1	Banan2
P-Erum 1-A + P-Erum 1-B	172	None	172	172	Multibands	172	172	172	Multibands
P-Erum 2-A + P-Erum 2-B	217	None	515	500 + 900	280+500+1200	500+900	500+900	500+900	500+1200
P-Erum 3-A + P-Erum 3-B	508	None	508	None	560	508+1900	508+1900	None	508
P-Erum 4-A + P-Erum 4-B	642	None	642	642	642 <sup>a</sup>	642	642	642	642
P-Erum 5-A + P-Erum 5-B	238	None	238	None	238	238	238	238	238
P-Erum 6-A + P-Erum 6-B	None	127	None	127	None	127	127	127	None
P-Erum 7-A + P-Erum 7-B	2791	2395	2791+ 500	2395	2395	2395	2395	2395	None
P-Erum 8-A + P-Erum 8-B	552+1071 +480	492	1200 + 500 <sup>a</sup>	492	492	492	492	492	552+1071 +480
P-Erum 9-A + P-Erum 9-B	1361	1178	1361	1000	1000	1000	1000	1000	1178+1361
P-Erum 10-A + P-Erum 10-B	1095+300	1691	820	820	820	820	820	820	1095

a: presence of additional multiple bands is observed

# EXAMPLE 5: DIFFERENTIAL SPECIFIC PCR DETECTION AND IDENTIFICATION OF ATTENUATED AND DIFFERING DERIVATES OF STRAINS GARDEL AND SENEGAL

The primers listed in Table 2 also allow the specific identification of attenuated variants of known strains of *E. ruminantium*.

The following variants were tested:

- attenuated derivatives of strains Gardel and Senegal denominated Gatt (for Gardel-attenuated) and Satt (for Senegal-attenuated), respectively. The strain Gatt was obtained from the virulent strain Gardel through 248 successive passages BUEC cells whereas strain Satt was obtained from the virulent strain Senegal following 64 passages on BUEC cells. Both the Gatt and Satt strains display an attenuated phenotype characterized by a lack of virulence.

- strain Gardel CTVM, which is a subset of strain Gardel maintained in a differing cell environment, and was reported has having undergone mutations in the *map1* operon and displaying a diverging phenotype (Bekker *et al.*, 2004).

DNA is extracted from elementary bodies of *E. ruminantium* and PCR amplification performed as described in Example 3.

The results are shown in Table 5, Figure 4, and Figure 5.

## Legend of Figure 4:

- A. PCR detection with primers P-Erum 7-A + P-Erum 7-B
- B. PCR detection with primers P-Erum 8-A + P-Erum 8-B
- C. PCR detection with primers P-Erum 9-A + P-Erum 9-B
- D. PCR detection with primers P-Erum 10-A + P-Erum 10-B

MW1: Molecular weight marker (100pb DNA ladder); MW2: Molecular weight marker (100pb DNA ladder); 1: Strain Bankouma; 2: Strain Bekuy; 3: Strain Lamba; 4: Strain Banan 1; 5: Strain Banan 2; 6: Strain Gardel attenuated (Gatt); 7: Strain Gardel CTVM; 8: Strain Senegal; 9: Strain Senegal attenuated (Satt); NC: Negative control; G. Strain Gardel; W: Strain Welgevonden.

## Legend of Figure 5:

- A. PCR analysis of virulent and attenuated strains with primers P-Erum 1-A + P-Erum 1-B  
MW1: Molecular weight marker (100-pb DNA ladder); MW2: Molecular weight marker (100pb DNA ladder); 1: Strain Gardel attenuated (Gatt); 2: Strain Gardel; 3: Negative control.
- B. PCR analysis of virulent and attenuated strains with primers P-Erum 2-A + P-Erum 2-B and P-Erum 6-A + P-Erum 6-B  
MW1: Molecular weight marker (100-pb DNA ladder); MW2: Molecular weight marker (100pb DNA ladder); Analysis with P-Erum 2-A + P-Erum 2-B of 1: Strain Gardel attenuated (Gatt); 2: Strain Gardel; 3: Negative control; Analysis with P-Erum 6-A + P-Erum 6-B of 4:

Strain Gardel attenuated (Gatt); 5: Strain Gardel; 6: Strain Welgevonden; 3: Negative control.

- C. PCR analysis of virulent and attenuated strains with primers P-Erum 3-A + P-Erum 3-B, P-Erum 4-A + P-Erum 4-B and P-Erum 5-A + P-Erum 5-B

- 5 MW1: Molecular weight marker (100-pb DNA ladder); MW2: Molecular weight marker (100-pb DNA ladder); Analysis with P-Erum 4-A + P-Erum 4-B of 1: Strain Gardel attenuated (Gatt); 2: Strain Gardel; 3: Negative control; Analysis with P-Erum 5-A + P-Erum 5-B of 4: Strain Gardel attenuated (Gatt); 5: Strain Gardel; 6: Negative control; Analysis with P-Erum 3-A + P-Erum 3-B of 7: Strain Gardel attenuated (Gatt); 8: Strain Gardel; 9: Strain Gardel;  
10 10: Negative control.

As shown in Table 5, Fig. 4 and Fig. 5, the combined use of all the pairs of primers described in Table 2 also allowed for the differential identification and discrimination of variants and attenuated derivatives of known strains. The PCR reactions results are summarized in Table 5. The overall PCR patterns generated on the strain Gardel and two of its derivatives, the strain Gardel CTVM and the attenuated strain Gatt show the presence of slight variations (Table 5, Fig. 4, Fig. 5). Furthermore, each strain is characterized by a specific pattern. The strain Gatt differs from the parental virulent strain Gardel by the products from primers pairs P-Erum 6-A + P-Erum 6-B and P-Erum 7-A + P-Erum 7-B, whereas the strain Gardel CTVM differs from the parental strain Gardel by the product from the primers pair P-Erum 6-A + P-Erum 6-B. Similarly, the primers pair P-Erum 7-A + P-Erum 7-B allows for discrimination between the strain Gardel CTVM and the attenuated Gardel strain Gatt. A similar situation is observed between the parental virulent strain Senegal and its attenuated derivative Satt (Table 5, Fig. 4). The virulent strain Senegal and the attenuated strains Satt differ by the PCR product from the primer pairs P-Erum 2-A + P-Erum 2-B, P-Erum 3-A + P-Erum 3-B, P-Erum 6-A + P-Erum 6-B and P-Erum 7-A + P-Erum 7-B.



Table 5: Differential identification of attenuated Gardel and Senegal strains of *E. ruminantium*

Primer combination	Strain				
	Gardel (Virulent)	Gatt (attenuated Gardel)	Gardel CTVM	Senegal	Satt (attenuated Senegal)
P-Erum 1-A + P-Erum 1-B	172	172	172	172	172
P-Erum 2-A + P-Erum 2-B	217	217	217	500 + 900	500 + 900+210
P-Erum 3-A + P-Erum 3-B	508	508	508	None	508+1900
P-Erum 4-A + P-Erum 4-B	642	642	642	642	642
P-Erum 5-A + P-Erum 5-B	238	238	238	None	238
P-Erum 6-A + P-Erum 6-B	None	127	127	127	127
P-Erum 7-A + P-Erum 7-B	2791	None	2791	2395	2395+700+300
P-Erum 8-A + P-Erum 8-B	552+1071+480	552+1071+480	552+1071+480	492	492
P-Erum 9-A + P-Erum 9-B	1361	1361	1361	1000	1000
P-Erum 10-A + P-Erum 10-B	1095+300	1095+300	1095+300	820	820

**EXAMPLE 6: ABSENCE OF CROSS-REACTION WITH OTHER RICKETTSIALES**

To verify the specificity assessment of the primers listed in Table 2, they were tested on *Rickettsiales* belonging to other species and genera than *E. ruminantium* i.e.

5 *Ehrlichia canis*, *Anaplasma platys* and *Anaplasma marginale*.

DNA extraction and PCR amplification were performed as described in Example 3.

The results are shown in Figures 6, 7, and 8.

Legend of Figure 6:

- 10 - A. PCR detection with probe EHR16S specific to *Ehrlichia spp.* 16S rDNA  
MW: Molecular weight marker (100-bp ladder); 1, 2 and 3: DNA isolated from blood samples from dogs infected with *Anaplasma platys*, 4, 5 and 8: DNA isolated from blood samples from dogs infected with *Ehrlichia canis*; 6: *Ehrlichia canis* positive control; 7: Negative control; 9 and 10: Control DNA isolated from blood samples from non-infected dogs.
- 15 - B. PCR detection with Nested PCR probes specific to *Anaplasma platys*  
MW: Molecular weight marker (100-bp ladder); 1, 2 and 3: DNA isolated from blood samples from dogs infected with *Anaplasma platys*.

- C. PCR detection with Nested PCR probes specific to *Ehrlichia canis*

MW: Molecular weight marker (100-bp ladder); 4, 5 and 8: DNA isolated from blood samples from dogs infected with *Ehrlichia canis*; 11: DNA from canine monocytes cultures infected with *E. canis* (supernatant); 12: DNA from canine monocytes cultures infected with *E. canis* (pellet).

Legend of Figure 7:

A. PCR detection with primers P-Erum 1-A + P-Erum 1-B

B. PCR detection with primers P-Erum 2-A + P-Erum 2-B

C. PCR detection with primers P-Erum 3-A + P-Erum 3-B

D. PCR detection with primers P-Erum 4-A + P-Erum 4-B

E. PCR detection with primers P-Erum 5-A + P-Erum 5-B

F. PCR detection with primers P-Erum 6-A + P-Erum 6-B

MW1: Molecular weight marker (100pb DNA ladder); MW2: Molecular weight marker (1 *HindIII/EcoRI*); 1, 2 and 3: DNA isolated from blood samples from dogs infected with *Anaplasma platys*, 4, 5 and 8: DNA isolated from blood samples from dogs infected with *Ehrlichia canis*; 9 and 10: Control DNA isolated from blood samples from non-infected dogs, 11: DNA from canine monocytes cultures infected with *E. canis* (supernatant); 12: DNA from canine monocytes cultures infected with *E. canis* (pellet); Am: DNA from *Anaplasma marginale*; G: DNA from strain Gardel; NC: Negative control.

Legend of Figure 8:

A. PCR detection with primers P-Erum 7-A + P-Erum 7-B

B. PCR detection with primers P-Erum 8-A + P-Erum 8-B

C. PCR detection with primers P-Erum 9-A + P-Erum 9-B

D. PCR detection with primers P-Erum 10-A + P-Erum 10-B

MW1: Molecular weight marker (100pb DNA ladder); MW2: Molecular weight marker (1 *HindIII/EcoRI*); 1, 2 and 3: DNA isolated from blood samples from dogs infected with *Anaplasma platys*, 4, 5 and 8: DNA isolated from blood samples from dogs infected with *Ehrlichia canis*; 9 and 10: Control DNA isolated from blood samples from non-infected dogs, 11: DNA from canine monocytes cultures infected with *E. canis* (supernatant); 12: DNA from canine monocytes cultures infected with *E. canis* (pellet); Am: DNA from *Anaplasma marginale*; G: DNA from strain Gardel; W: DNA from strain Welgevonden; NC: Negative control.

Fig. 6 indicates that the samples used indeed contain DNA from *A. platys* and *E. canis* as demonstrated by their recognition by 16S rDNA-specific primers and primers for nested PCR specific to each species. As shown in Fig. 7 and Fig. 8, the pairs of primers described in Table 2 are strictly specific to *E. ruminantium* and display no cross-reaction with other related *Rickettsiales* since no specific PCR product could be detected on *E. canis*, *A. platys* and *A. marginale* (Fig. 7 and Fig.8). Whereas no PCR products are detectable on

whatever pair of primers was used, PCR products were visible on *A. platys* and *E. canis*. However, all these PCR products are generated by cross-reactions with canine blood cells as shown by the detection of these same bands on non-infected canine cells (Fig. 7 and Fig. 8). This demonstrate that the primers described in Table 2 and targeting the target genes  
5 described in Table 1 allow for specific identification of *E. ruminantium* and discrimination between strains of *E. ruminantium* even when other *Rickettsiales* are present.

The tools provided by the invention allow thus both for specific detection of *E. ruminantium*, even in presence of contaminating related *Rickettsiales*, for specific discrimination between different strains of *E. ruminantium* and for specific discrimination  
10 between a virulent strains and its vaccinal attenuated derivates. This in turn allows for monitoring of vaccination.

## CLAIMS

1) Use of the following set of target genes :

Erum1, defined by the sequence SEQ ID NO: 6

Erum2, defined by the sequence SEQ ID NO: 3

Erum3, defined by the sequence SEQ ID NO: 1

Erum4, defined by the sequence SEQ ID NO: 4

Erum5, defined by the sequence SEQ ID NO: 2

Erum6, defined by the sequence SEQ ID NO: 5

Erum7, defined by the sequence SEQ ID NO: 13

Erum8, defined by the sequence SEQ ID NO: 15

Erum9, defined by the sequence SEQ ID NO: 14

Erum10, defined by the sequence SEQ ID NO: 8,

for the strain-specific detection of *Ehrlichia ruminantium*.

2) A method for the strain-specific detection of *Ehrlichia ruminantium* wherein said method comprises detecting, for each of the genes Erum1 to Erum10, whether an allele of said gene is present in the bacteria to be tested, and determining the form of said allele.

3) A method of claim 2, which comprises performing PCR amplification of all the target genes Erum 1 to Erum10, and checking, for each of these genes, the presence of one or more amplification product(s), and the size of said amplification product(s).




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5






Figure 6




Figure 7




Figure 8

## SEQUENCE LISTING

<110> - CENTRE DE COOPERATION INTERNATIONALE EN RECHERCHE  
 AGRONOMIQUE POUR LE DEVELOPPEMENT (CIRAD)  
 - CENTRE NATIONAL DE LA RECHERCHE SCIENTIFIQUE  
 - FRUTOS, Roger  
 - VACHIER, Nathalie  
 - LEFRANCOIS, Thierry  
 - FERRAZ, Conception  
 - DEMAILLE, Jacques  
 - MARTINEZ, Dominique

<120> TARGET GENES FOR STRAIN-SPECIFIC DIAGNOSTIC OF EHRLICHIA  
 RUMINANTIIUM AND USE THEREOF

<130> MJP/mad-F1367-12/WO

<160> 35

<170> PatentIn version 3.3

<210> 1

<211> 630

<212> DNA

<213> Ehrlichia ruminantium

<400> 1

```
atgaaaggat ctttatctgc taaagttatt tctgaaaatc taccattagt agagatggaa      60
aaagcagttc ttagtcctac tgctcgtatt tttctcacta atcataagtt gggacctgtc     120
atggaccttg gaatttatat ctttaatacat catagtaatc ttcgtttatt aacgaaggaa     180
aacctttatc ctgctaataa cctaagtaaa attggtaaag tgggtgctttg taaacctttg     240
tctataggca atggcataca tacagtacat atgtacttta atgaactcga agcttttaaaa     300
gaattcggag gattagaaaa tgctcgtctt acaacagtac gtccggactc ccccttgcac     360
acacatacat ctaaaaaaaaa gaaatcatta ttacaaaac gttcagatac ttgctataca     420
ctattatgtg aggaatctta tacagatcca aataataccg aaactgatag tacagtaaaa     480
gcaatatcac ataatgaaga agaagaaggt gcagtaagag gagatatacc acaatatcaa     540
ctttccaatg ccgaagcact aggtcgtggt cttgcttttt tccatgatgt tgcaagtaat     600
tttgaaacat tatgcagaag ataccattaa      630
```

<210> 2

<211> 303

<212> DNA

<213> Ehrlichia ruminantium

<400> 2

```
atggatttaa ataaactaat aaagagatta gtattttcat ttgtaatgat taattttggt      60
aataggtttt ttagtaatac agaaagtgaag agcttgcatc taagtgatag ttacgacac     120
tattattatt ttctatgttt gtgccatgca gtaatggggg ttattatagt aaatacagat     180
```

ggatataaca tccttgagga ttttatgttc tcagaacaaa tcgtaggtag agaaaatgca 240  
 gaaatgcttt caatatcaga tacagagggg ggggggggag agcttagtag aagaaaattc 300  
 tag 303

<210> 3  
 <211> 270  
 <212> DNA  
 <213> Ehrlichia ruminantium

<400> 3  
 atgtatttag tctatttagt agctggtttt gtggtactat atagtaatta tcgagatata 60  
 aattatgata aaaaacttgc tattctttat tctaggggag aagatgatga atataaatat 120  
 gttcctagga aagagcagaa taatcaatat tattttcata taaaattgta tagtgtaag 180  
 ttaaatttaa tgtcagatat tgttcaatta gatgttataa tgtaaaaagg attttattat 240  
 agcaatatgt ttaatgtctt tttatttttaa 270

<210> 4  
 <211> 828  
 <212> DNA  
 <213> Ehrlichia ruminantium

<400> 4  
 gtgtgttact taattggtaa ttttatgtta ttcaaataca atcctcaaaa tactaaagaa 60  
 ttacatgatg cagctttaaa ttgtttacgt catacaagat tatatgcata tagctaccgt 120  
 tgtataggac atactgaacc taatggaaca ctacatgtat tcataagtaa agataaatca 180  
 aataatttgt gtttaccaaa agaagggtat tctctattct atatagaatg tagtctatct 240  
 gataagagag tatctcagaa tcaggaaata agagatatga tgcaagcagt tgtccgccac 300  
 aaaattaacc gccttgcttt taataaccct cacacgacac ctacataga tgtaggcatt 360  
 tatatttttaa taaataaaag taaccttaat atgttaacaa aagaacatat aacacctacc 420  
 aacaacatgg acagtgttgg ccatatgata ttatgcaaac ctgtacgtgc agctaattgt 480  
 ttactctcat tagacttctt attcaatgaa gaagaagctt taaaagagct tggaggatta 540  
 caaaatgcag tattttacgat aatagaaact acaccaccta ttaccaaaaa atcattattc 600  
 agaagacatt cactgggtta ttcacaacta tcagaagaac atagtaaacc tgaaacaatt 660  
 accagtagta ctattacaga gagtataaca agagaagaag cacaatcaag taaacaagag 720  
 gaaggattag aaacacatca gctttccacc aatgtagtaa cacatggat caattattta 780  
 actaatgtct cacttgcttt tgaacagcta tgtacaaaat atcattaa 828

<210> 5

<211> 225  
 <212> DNA  
 <213> Ehrlichia ruminantium

<400> 5  
 ttgtacatag tatgtcttta tataaaagta agaattgatg attctactga tgttattact 60  
 tataactcta aaaaaaatat gtgtaaatta caattaactc agaaaaagaa tagatcattt 120  
 atatatattgg ttaacagata ctatcataaa tcagaatata ggcttaccac actttcagtt 180  
 caatcttata gcaaattaga gcaactttat aacaatatcc agtaa 225

<210> 6  
 <211> 186  
 <212> DNA  
 <213> Ehrlichia ruminantium

<400> 6  
 atgagtcaca gttttattga gtttaaacia atcaattatt acgatattaa cgcaatatat 60  
 acaatatcat ttgtaacaca tatcaataat ttatatccaa aatataagag aaaaattatt 120  
 ataactctgc ttaatacact aattaacaat attacttctt gtgattttga gtgtaataaa 180  
 caatga 186

<210> 7  
 <211> 3522  
 <212> DNA  
 <213> Ehrlichia ruminantium

<400> 7  
 atgaatgaga taatcctata cacagcagtg tcgctgtttt ttatatgtgt ttactatggt 60  
 ctgcttgtgg ttaggtttgt atgttatgtg ttgagtgtta tgaagtataa gtcaaaggaa 120  
 ttggacatat cagataatta tacaaaaagt agggttactt attgtagtca gagtgaatat 180  
 gaaaagtacg aaatggacac tttatctgga aaagatggta ttgaatttct aaaatcagtt 240  
 taccataatg atagtgatga tataggatcat gttttaaaat caaaatctac tgtttcatct 300  
 accaaaatgg atcaggtaac acatcaagtt cctggcggtc aaactataga acacgatagt 360  
 gcgatagaag gtcaccaagt tatggataag gaaaatgctg gtgttggtgt tcaactatagt 420  
 catactgaaa ctactataaa aacaagtctt agttttaaat ctgatgttat gggtgatact 480  
 aaggataaat ctgtagagaa aaaagtagta cctgaaaata ctataagaat aaatgaaaaa 540  
 aagagagatg tttttgtaag tgctagtatt caaactgata taaaaagtaa tcaagttaaa 600  
 ttatctagtt ctgtattaga aaaaccagat gagaaaagtg atgttactga tacagcgtgt 660  
 acaggtagta ctaaggataa atctgtagag gaaaaagtag tacctgaagg tgatactata 720  
 agaataaatg aaaaaaagag agatgttttt gtaagtgtca gtgctcaaac tggatgatgt 780

aaaagtgatc aagttaaatt atctggttct agattagaaa aactagatga gagaaaggat	840
gttactgata caggttgtgc aggtagtact aaggacaaat ctgtagagaa aaaagtagta	900
tctgaaggta ctgctataag agatgaaaag gagagtagtg ttgctagaag tgttggtgtt	960
acttttaatc ttcaaagtgg taatgtaaaa gatgataaag taaaactatc aggtgtagat	1020
ttaggtaaaa tagaggattc agttttatct gcttctagtt gtgaaactac tgttaaggat	1080
aataagcctg ttatatgtgt tggaaaagaa agtacgtttc aattagcttc aagtttgat	1140
ttggttaata ctgttgaaga tagttcaaga aatactcgtg gtttaagtga aacttgttct	1200
ttaatgttag attttgacag aaatggtaat cctgtatacg aagaggcaac tagtaagta	1260
gtgcctagtt tctatcctga taatgttata tatcacacta aagaaaaaca ttgtggtgtt	1320
gatcttcctc aatcagaaga tcaactttat tcatgtatta ctaatgtgca tagtcaatat	1380
gatgtgactg aaaatagtgt aagtgtatat ccgcgtgatt tggttcctga tgatataaaa	1440
caagctaaac agaatgaaga tactaaacag ggtgctttta tagctacagg ttctacaacc	1500
gcggtcgcgc atagtcaata tgatgtgact gaaaatagcg taagtgtatg tcagagtgat	1560
ttggttcctg atgatataaa acaagctaaa cagagtgaag atactaaaca gggtgctttt	1620
atagctacag gttctacaac cgcggtcgc catagtcaat atgatgtgac tgaaaatagt	1680
gttagtgtat atcagagtga cttagtttct gataatataa aacaagctaa acagaatgaa	1740
gatactaagc agggtgcttt tatagctaca ggttctacaa ccgcggtcgc gcatagtcaa	1800
tatgatatga ctgaaaatag cgtaagtgtg tgtcagagtg acttagttcc tgatggtgta	1860
aaacaatcta aacagcatga agatactaag cagggtgctt ttatagttac aggttctgta	1920
tctgctaagt tagatattgt tgatgtagtt agtttagggg aaaaacgtga tattgatgaa	1980
aaagttgtta agtcatcagg ttgtactact gctgattcag ttagtaatcc tgtaggtatg	2040
gataaagttc aatattgtgt acctgactta gagatgagag taaaaatgga tcttgtagaa	2100
gatcaccata atatggctag tatggaaaaa tgttatcctg atagagaagt tgttgagcaa	2160
ttaagtaatg ttactacttg tttggttagt actccagtaa ttgaacatag agttcatagt	2220
gttgagtctg ttgcagagtt acaagtaaaa ataggtcctt tagatgaggg aaaatgtaaa	2280
gacagtgtgg taaggagctc atcatttact agtgatacat gtttaaaaga tacagggtgca	2340
acaatgactg tagaagaata tggtataaaa cctagtacag gtctttgtgc tagtaggggt	2400
gatgatagtg tttcttctat gattggtata ggttcgtatt ttatagataa gatgatttgt	2460
gatattgata ctactgtgca gcttaataat acattttcta ctttagaaaa aagaaaaaac	2520
tgttttatag ataatatata aaaaaataat gaaaaaatat ttagtaacct tgttaatat	2580
atggatttaa taaaagaaac ggtaggtatt caattttttg atactaaaag tacagatgat	2640

atatccaggt atgtaatgga acaatctagt ggtgtttatg atgatgttat gtcacaaatg 2700  
 cttatccaag atgaaaaata tttatttaag gtctttaaac atattattcc gggttttgcct 2760  
 aaaatattct ttaacaatga tcctatatct tcaatggaat ggaaattagt agatgaattg 2820  
 ttctctatga gaagggcagt cttacaagat aatgtgtatt ttcaaaggat attttattgt 2880  
 atagtgtgtg catgtgaaaa aactgcaggt acaataaaga aaattcagtc gttatctaaa 2940  
 cagtgtgatg aaatacgaga aaagattaaa aagtgtaatc taaggcaagg aaagaagaaa 3000  
 agtgcattgt cgaaatttac agatcatttt agtgaaaaaa aggaagacct gttgtgttta 3060  
 ttagataaaa tagaaaaaga actgaattta actaagcaag ttacactaa tcttatagca 3120  
 gaaaaagagg cgttattaac aggagatgtt gcttatataa gatattttgt atcacgtatt 3180  
 gtttttgata gttggaaatt tgatgataag gctaaacagg ttgtcaaaaa tataaagaac 3240  
 ctagcaccat atgtgttatg tgatgtgttg tatgaagaag aaaaaaata tctagggttg 3300  
 gtgaagtgtg ttgtttgtga gtacacggtt ttttataaag atatagataa ttttttacct 3360  
 atagttcaac aatatcatga tcgacgacaa tctagaagtg ctgcagccca aaaattttat 3420  
 gatcaggaaa ttgatggtgt tcttcctatg gatactttag aagggttagg ggatcttgta 3480  
 gctatggaat taggacaaaa cagtaaattgt aatgcacatt aa 3522

<210> 8  
 <211> 4122  
 <212> DNA  
 <213> Ehrlichia ruminantium

<400> 8  
 atgaatgaga taatcctata cacagcagta tcaactgtttt ttatatgtat ttactatgtt 60  
 ctgcttgtgg ctaggtttgt gtgttatgtg ttaagtatta tgaagtataa gtcaagagaa 120  
 ttggatatat cggataatga tacaaaaagt agggttactt attgtagtca gagtgagtat 180  
 gaatatggaa agtacgagat ggaaacttta tctggaaaag atggattga atttctaaaa 240  
 tcagtttacc atagtgatag tgatgatgta ggtgatgttt taaaatcaaa atctactgtc 300  
 tcatctacca aaatggatca ggtaacacat caaatttctg acgttcaaac tatagaacgc 360  
 gataatgtag aagggtcaaca agttatggtt aaggaaaatg ctggtgttgg tgttcactat 420  
 aatcactactg aaactattat aaaaacaagt cttagtttta aatctgatgt tatggttgat 480  
 actaaggata aatctataga ggaaaaagta gtacctgaag gtgatactat aagaataaat 540  
 gaaaaaaaga gagatgtttt tgtaagtgtc agtgctcaaa ctgatatgaa aagtaatcaa 600  
 gttagattat ctggttctag attagagaaa ccagatgaga gaagggtgt tactgataca 660  
 gcgtgtacag gtagtactaa ggataaatct gtagaggaaa aagtagtacc tgaagggtgat 720



actataagaa taaatgaaaa aaagagagat gtttttgttaa gtgctagtgc tcaaactgat	780
atgaaaagta atcaagtttag attatctggt tctagattag agaaaccaga tgagaaaagg	840
gatgttactg atacagcgtg tacaggtagt actaaggata aatctataga ggaaaaagta	900
gtacctgaag gtgatactat aagaataaat gaaaaaaaga gagatgtttt tgtaagtgc	960
agtgcctcaa ctggtgatat gaaaagtgat cacattaaat tatctggttc tagattagag	1020
aaaccagatg agaaaagga tggtactgat acagcgtgta caggtagtac taaggataaa	1080
tctgtagagg aaaaagtagt acctgaagg gatactataa gaataaatga aaaaaagaga	1140
gatgtttttg taagtgcctag tgctcaaact ggtgatatga aaagtgatca cattaaatta	1200
tctggttcta gattagagaa accagatgag agaagggatg ttactgatac aggttgctacg	1260
ggtaatacta aggataaatc tgtagaggaa aaagtagtac ctgaagggtga tactataaga	1320
ataaatgaaa aaaagagaga tgtttttgta agtgctagtg ctcaaactgg tgatatgaaa	1380
agtaatcaag ttaaattatc tggttctaga ttagaaaaac tagatgagag aaaggatgtt	1440
actgatacag gttgtacggg taataactaag gataaatctg tagagaaaaa agtagtatct	1500
gaaggctact ctataagaga tgaaggag agtagtggtg ctagaagtgt tgatgctact	1560
tttaatcttc aaagtggtaa tgtaaaagat gataaagtaa aactatcagg tgtagattta	1620
ggtaaaatag aggattcagt tttatctgct tctagttgtg aaactactgt taaggataat	1680
aagcctgtta tatgtgttg aaaagaaagt acgtttcaat tagcttcaag tttggatttg	1740
gttaatgctg ttgaagatag ttcaagaaat acttggtggtt taagtgaac ttgttcttta	1800
atgttagatt ttgacagaaa tggtaatcct gtatacgaag aggcaactag taagttagtg	1860
cctagtttct atcctgataa tggtatatat cacactaaag aaaaacattg tgggtgtgat	1920
cttctcaat cagaagatca actttattca tgtattacta atgtgcatag tcaatatgat	1980
gtgactgaaa atagtgtgaa tgtatatccg cgtgatttg ttcctgatga tataaaacaa	2040
gctaaacaga atgaagatac taaacagggt gcttttatag ctacagggtc tacaaccgcg	2100
gctgcgcata gtcaatatga tgtgactgaa aatagcgtaa gtgtatgtca gagtgactta	2160
gttctctgat atataaaaca agctaaacag aatgaagata ctaaacaggg tgcttttata	2220
gctacagggt ctacaaccgc ggctgcgc atgtcaatatg atgtgactga aaatagtgtt	2280
agtgatatat agagtgactt agttcctgat gatataaaac aagctaaaca gaatgaagat	2340
actaagcagg gtgcttttat agctacagg tctgcaaccg cggctgcgca tagtcaatat	2400
gatatgactg aaaatagcgt aagtgtatgt cagagtgatt tggttcctga tgatataaaa	2460
caagctaaac agaataaga tactaagcag ggtgctttta tagttacagg ttctgtatct	2520

```

gctaagttag atattgttga thtagttaat ttaggggaaa aacgtgatat tgatgaaaaa 2580
gttggttaagt catcaggttg tactactgct gattcagtta gtaatcctgt aggtatggat 2640
aaagttcaat attgtgtacc tgacttagag aggagagtga aaatggatct tgtagaagat 2700
cactataata tggctagtat ggaaaaatgt tatcctgata gagaagttgt tgagcaatta 2760
agtaatgtta ctacttgttt ggtagtagt ccagtaattg agcatagagt tcatagtgtt 2820
gagtctgttg cagagttaca agtaaaaaata ggtccttttag atgagggaaa atgtagagac 2880
agtgtggtaa tgagctcatc atttactagt gatacatgtt taaaagatac aggtgcaaca 2940
atgactgtag aagaatatgg taataaacct agtacaggtc tttgtgctag taggggtgat 3000
gatagtgttt cttctatgat tggtaggggt tcgtatttta tagataagat gatttgtgat 3060
attgatacta ctgtgcagct taataatata ttttctactt tagaaaaaag aaaaaaacat 3120
tttatagatg atattaaaaa aaataatgaa aaaatattta gtaaccttgt taatattatg 3180
gatttaataa aagaaacggg aggtattcaa ttttttgata ctaaaagtac agatgatata 3240
tccaggtagt taatggaaca atctagtggg gtttatgatg atgttatgtc acaaatgctt 3300
atccaagatg aaaaatattt atttaaggtc tttaaacata ttattccggg ttttgctaaa 3360
atattcttta acaatgatcc tatatcttca atggaatgga aattagtaga tgaattgttc 3420
tctatgagaa gggcagtcctt acaagataat gtgtattttc aaaggatatt ttattgtata 3480
gtgtgtgcat gtgaaaaaac tgcagggtgca ataaagaaaa ttcagtcatt atctaaacag 3540
tgtgatgaaa tacgagaaaa gattaaaaag tgtaatctaa ggcaaggaaa gaagaaaagt 3600
gcattgtcga aatttacaga tcatttttagt gaaaaaaagg aagacctgtt gtgtttatta 3660
gataaaatag aaaaagaact gaatttaact aagcaagttt acactaatct tatagcagaa 3720
aaagaggcgt tattaacagg agatgttgct tatataagat attttgtatc acgtattgtt 3780
tttgatagtt ggaaatttga tgataaggct aaacaagtta tcaaaaatat aaagaaccta 3840
gcaccatagt tgttacgtga tgtgttgat gaagaggaaa aaaaatatct aggttttggtg 3900
aagtgtattg tttgtgagta cacggttttt tataaagata tagatgattt tttacctgcg 3960
gttcaagaat atcataatcg acgacaatct agaagtgtg cagcccgaat attttatgat 4020
caggaaattg atggatttct tcttcctatg gatactttag aagatgtagg ggatcttgta 4080
gctatggaat taggacagaa cagtaaatgt aatgcacatt aa 4122

```

&lt;210&gt; 9

&lt;211&gt; 2856

&lt;212&gt; DNA

&lt;213&gt; Ehrlichia ruminantium

&lt;400&gt; 9

ttgcataaaa tcatgcttac atcacttaaa actacagtta ctgataataa actaagatct	60
agtattatta atggatctag tgttaatttt tttaaaaaag gcaacattat tttttctgta	120
tattatacaa gaaataatgt taatggatat gatgtaatat gtgagattca acatgggtgct	180
tctattttatt atatgaaaat taatgatcat gcaattattg atcgtcggac aacaaattat	240
ccaccagaga tgttatttgt aaagaatagt aatgatgatt tgatatttat tgttattcct	300
gaagaaagta aaggctgtaa agcttttagtt atcaaaatat ataagataaa ttataatcct	360
aatgtgtctt tatccaaatt acatgactta caattaatta gttgcaataa ctatctcaaa	420
gaagaagtga aatatcctat tattttacat caggatacgg gtggtaggat tgttggttatt	480
gcaagagtag ataatgacta tcgaggatgat gttacagata gattgtatgt gatgtggcaa	540
ttgagatata ataataatag atttgaaatt atagggttaa gtaatgggta taggcaattt	600
aatgctgact atttatttaa gcattctggt tatattaatc gtggtaaatg tcatgataga	660
ctaattgtta aattaggatc ggatagtttt ataaattttc tttatgttgg gaaacatatt	720
tctagagatt acaatttttt ttctagtata tatgatttat ctataaatta taatatgcat	780
cttgatccag aagaatgttt ggtgggttct ttttatgggt gtaatgctag tagtggtagg	840
aaatataata ttccataatag ctgtattcat gttattgata tttttcgtga tgatgggaat	900
gtatatatag catatattgg tactgtattt aatagtacat ttaagaataa aaagcagttg	960
gttattgttt atactatggg tgatgaacaa tcgcctgtgt atgattttat gcagattaca	1020
gaagatatta gtgccatata tataaattct actgaaaata ttttagcaat aacgactatg	1080
ggaagtgatt atcttgtaaa atatgagatt tcaaaattac agttaaaatt agggattgtt	1140
gatcatgttg atgttatataa aattccacgt aatgtagtga aaaatattgc taattttaca	1200
tatgttggtg atacagtttt aggggttgat agtggtgaac atattaatat tcgtaatgta	1260
ttagctacaa aatcaactgt taatgataaa gtatctcagt tttttttaa tattagagaa	1320
atggaatttg gtgatttatt taagagttgg agtggtgaat ataatgactt gttaataggt	1380
tatactatgc ctgctagtta tgggtgaaat tatactacag aatatttaag cgatgttata	1440
actgtttcag gtaatgcagg ttttgtagag aagttcatat caactagtaa aatgggtgat	1500
gtatttaaga ttacagataa cttaattaat tatactagtg taaaccctac taatcatatg	1560
gcacatgtga cattgcaatc aaaattatca gatgggtgagg gtattacaga gcgtgcgggt	1620
aatagatcag ataattctgt aagtgaaggt ttagctacag gattggttct tactagtaaa	1680
agtgatgatt tgtttaaaag tacggctagt cctattaatc atgcttttgg ttatgtaata	1740
aagcctacta tccacgtaac gcatgttaca ttgcaaccga agtcaccata tggtaaagag	1800
gttgtgaggc atataaatcc taaaacggat aattctatac atacaagttc aataccaaga	1860

tcaataactga ctaatagaag cgatggtgta tttaaaagta cagctagtcc tattaatcat	1920
gcttttggtt atgtaataaa gcctactcgc catgtaacgc atgttacatt gcaaccgaag	1980
tcaccatattg gtaaagaggt tgtgaggcat ataaatccta aaacggataa ttctatacat	2040
acaagttcaa taccaagatc agtactgact agtaaaagct atgatgtatt taaaagtacg	2100
gctagtccta ttaatcatgc ttttggttat atgaaacctg ctagttctgt tgtagtgcc	2160
ttaggtgata ctgatgtttc aaagcaagtg gaaagtgtta gtaatgttcc agtacatctt	2220
actcctacag taagatcagt attagtaggt gatgcgtatc atgtatctgg cagtgaaaaa	2280
gatagtgttg gacatgaaca agatttgggt catggtgatg ttagtactga tgttgattg	2340
aaactaatga gtgataatgt atccaacaat attagtaggc atgtgaataa ttcttttagct	2400
ataaaacata agatattagg tagaaaagta aagtataata taaggcgtag tactgttaga	2460
tctggtgtta atattcgcaa taaaagtaca gtgagtacaa gatatacatc tcatggcata	2520
caagaggcta ataatatgaa tgttacattg tttaatccta cacagcataa tattagtagt	2580
tataatggta gtttattaaa tagtaattct gcttttaata ctgaaagttc tgttgattat	2640
aaagtagtaa ttgcagtaat atctagtata ctgcttatct ttttattatt aggtggattt	2700
aaatgtataa agtggtatct agcaaagttg aatagaagaa ggatgtctaa taatgaacag	2760
ggatttgtga tttttaattt ggatagtatt caaagtagtg tttctggtgt gcaagtgaca	2820
aaaggtacca catctcgaat agagagtcta ttctag	2856

<210> 10  
 <211> 1659  
 <212> DNA  
 <213> Ehrlichia ruminantium

<400> 10	
atgttattca aaccoggttt acccatttcc aagattacag aatctctgca tagatcagtt	60
atacgtgagt tgaatagagc gtcagaaata catgttaata cgtgtcattg tataggagct	120
acaataaata aaaaaactct taatatctgc gttgataata agccaggtaa tcggtgtact	180
ccagtaggaa catctttatt tcgtatggaa tgtattatac ccgctcctgt aataaataat	240
ccacgtaata tatctttaca aaagttgaca caagtattgt ctagtccttt ttttaataaca	300
ctagaaccac ttaaggttga tgcataattt attgtaccag aggaagaatt aaagaatttc	360
atagatttag taaaaccttt atctagtatg ccacgtgaag gacttttacc tatttataat	420
attggtaa at tcggaacatt ttccttatgt aggccaaaaa gtataggtaa tagagatgta	480
aggcatgatg taccatttga cgaattcaaa gcttttaata ttttaggagg tctagaggat	540
tctatttttt ttaaaacacc ctcatcaatc cctgatatca caaaacgtaa taaaaacaa	600

```

agcatagcag atagtaaaca aaaaaaagtt gtagtaactg gtaaggaaat tcaacacaaa 660
atacaacata taaaaaaaaat gttttctaga gtttctacta cacaatgttc accttcaagt 720
acaccagtca gtgctcaaat gacacataat atagaagaaa aaacagcaag tagtccgcaa 780
aagccagcta tccaaaaggt tatagtaact agcaacaac ctogtaaaga agaaatacaa 840
tttatatata caaagtttcc tgaggctcca gaagaacatt catcttcaaa tcaaacacaa 900
gagacaacaa gcaagcattt agaacagcat ctatcaaaaa gtatagtagg tgggtgcgcca 960
cttatacaaa aaggtacagt agatcctcaa caagttgtca gaccaaaaaac atttgcatct 1020
agtccttttt ataaagaatc gacacttcca actacaaaat atccatcttc aaatcaaaca 1080
caagaaacag caagcaagca tttagaacag catccatcaa aaattacaca aaaaggtaca 1140
ataaaccttc aacaagttgt tagaccaaaa acacaatttt catctagtcc tttttataaa 1200
gaacaggtag ttccaaatat aaaacatcca tcttcaaadc aaagacaaga gacagcaaac 1260
aagcatttag aacagcgtac attaaaaaaa agtacactag gtagcatgcc gccatctata 1320
caaaaaggta caatagatcc tcaacaagtt gttaggccaa aaacacaatc tgcatctagt 1380
cctttttaca aagaatcgac acttccaact acaaaacatc aaatgttaag tgttatagaa 1440
gaatcgacaa atagtagtgt accaattaat acattaagtt ctgaagaaat accacggttt 1500
ttcagtgtag attattttag tagttataaa gtattgtacg atacttacaaga ataatcttat 1560
aaagttgata ctttaccaac agcacctctc gtcccatcat gtcaacttga agatgatgta 1620
ttgttgaaa atagtaatcc ccatgtatct ttgaattaa 1659

```

```

<210> 11
<211> 873
<212> DNA
<213> Ehrlichia ruminantium

```

```

<400> 11
atgccactta ctttcgatct atatgcacat gaaaaaagat taaatcttct tctatgcaat 60
gctataaatt ccgctgctaa tctagttaat gaaatagata tatcttgtgt aggatatact 120
gatgaaaccg gtaaattagt ggcttttatt gaccctaaca ttccactaaa cttattccct 180
attcctcaaa atctatcctt atttcgtata agtgggtacta taccaattac cattataaca 240
aattctaatt ctcaagaatt atctaaagag tttgttttca cagaagggtga aataaatagt 300
ggccttgta agtgtgaaat gtattgttta gtaggtaatg aaaatcttga tgattttact 360
gaaatatgta gtaatcctaa aataggatat gaaaatttaa taaaatttc caataatatt 420
tacacacaat tatttacaga agatatatta aaatttccta ttaacataga acatgctcta 480
tcaaatatag caaattttaa tgagaatat atatatgcat ctgatctagt aaggaaagaa 540

```

aaacttaagc agcttaaaga aagaaatgat gatttatgta atgcaatatg ccatgcatgt 600  
 aatgaggaga atgtaacttc tgtaaaatgc ataggacata ctcttgacag taatcaactc 660  
 acagttcata ttaaatgtcc agaaagcctt ttacctatac ctcaaagtaa ctctctatctt 720  
 cttgtcgaga tgagtatatatt acctaatgtt atagggggca atcaaataatt atcctctact 780  
 tttgaattaa cagaaagtga atgtaaaaaa ggtgctctca attgtacaat gtactgttta 840  
 gttagtaagg agaaacttaa aaaattttac tga 873

<210> 12  
 <211> 1740  
 <212> DNA  
 <213> Ehrlichia ruminantium

<400> 12  
 atggctatac cacacactat tatagcaa atgccaat tatcttctac ttttaaagta 60  
 gtaaaaaatg atcttgggtat tgatagtgtat tatatcttat gcactgcata ctgtctagta 120  
 actaagcata atcttcaaga ttttactaat gaagtgtcat cggatgatgt aatagatgat 180  
 actacacaac aaaaacgtca aaaattttaa aatatcatta aattatgttc tgtaaaatgt 240  
 gtaacattac atacacaaga aatgttatca ttaaaccatta gtgaagaaga actaatcaac 300  
 gatatagggt tatgtaatgc aactttcaaa tttttaagta atctgcatca agaaaagatt 360  
 gatatactta gacaagcaaa taataaatta tgtcgtgaaa tatgtcttaa acttaataaa 420  
 catcaaaca actatataag atgtatagga aatactgttg ataatacaatt aatagttacc 480  
 actcagtgtc cacgagatct tcttcctttt cctaaaaatc aatctttatt cattataagg 540  
 ataaatatac cacctaacat tatattacac agtaaaatc taaataatac attttaaatta 600  
 acaaaaagtg aaaaagaaga ttattacatt aaatgtgata tgtattgcct agtatatgaa 660  
 gaagatatta agagtcttat tgatgtatgt gatgatttag ataagccata tattgaagag 720  
 ttaattcaac attgttctgt aaaatgtata aaattgtata cacaagaaat gttatcatta 780  
 aacattagtg aagaagaact aatcaacgat ataggggtat gtaatgcaga gttttaaata 840  
 gttgaaagta gacagataat tgaatcagta ctagacacat ttgagtatat cgaaatacaa 900  
 gcaaataaac tcctatgcag aattttgcct acactttgtg ctttatataa aaaagatttt 960  
 ctcatcaaaa agatacgttg tataggtaat actatagatc ctgaacaagg attaacaatt 1020  
 tatctctcta gtatatcttc aaaggaacac ttaccaactg ccaaaggtag atctttatctt 1080  
 ttaatacgaa gtaggatgtt aactgaagtt atattaagta ctctgaact agtgaatgta 1140  
 cataatctaa gtgatgaaga aatgtcgagt aagtatttat tatgtgatat atattgccta 1200  
 gtagataatc aaaacattaa tctatttaag aatctttgta caaagacaag acagttttct 1260

gatgtttgtaa ttacatgtga tgtaagatat attaggatat atacaaaaga tgctagaaaa 1320  
ttcccattta atgaggcaag tgtattaaaa caattaggaa atataaaagg aaaatatctc 1380  
aatgaacaag acttttaaagc attagttagt tctggacttt atactaaatc agcaagtga 1440  
tcttcatcag cagtatcaac tgaagaagaa tcaattatac aacaagaact ccatgtaaaa 1500  
cagagtttaa aatcaagatt atcacaaata agaaaacaac taacacctga ttcttcatca 1560  
tctaatacag tatcaagtga agatgatata gatacaccag cagaaattaa aagaaaaaga 1620  
gaagcaagac gtttaaaatt agcaaaatta caacaagaag aatcacaaac aacaggaata 1680  
ggtatgttgt ctgatcaaga agtttcacat cataaatctc aaagtaaaga tttagattag 1740

<210> 13  
<211> 3252  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 13  
ttgcataaaa ttatgcctac atcacttaaa accatagtta ctgatagtaa actaagatct 60  
agtattattg atggatctag tgttaatttt tttaaaaaag gtaacattat tttttctgta 120  
tattatacaa gaaataatgt tgatggatat gatgtaatat gtgagattca acatgggtgt 180  
tctattttatt atatgaaaat taatgatcat gcaattattg atcgtcgggc aacgaattat 240  
ccaccagaga tgttatttgt aaagaatagt aatgatgatt tgatatttat tgttattcct 300  
gaagaaagta aaggtagtaa agcttttagt atcaaaatat ataagataaa ttataatcct 360  
aatgtgtctt tatoccaatt acatgattta caattaatta gttgcaataa ctatctcaaa 420  
gaagaagtga aatatcctgt tattttacat caggatacgg ttggtaggat tgttggtatt 480  
gcaagagtag ataatgacta tcgaggtgat gttacagata gattgtatgt gatgtggcaa 540  
ttgagatatc ataatagtag atttgaaatt ataggtttaa gtaatgggta tagacgattt 600  
aatgctgcct acttatttta gcaattctgg tattaatac gtggaaaatg tcatgataga 660  
ctaattgtta aattaggatc ggatagtttt ataaattttc tttatgttgg gaaacatatt 720  
tctagagatt acaatttttt ttctagtata tatgatttat ctataaatta taatatgcat 780  
cttaatccag aagaatgttt ggtgggttct ttttatggtt gtaatgctag tagtggtagg 840  
agatataata ttccataatag ctgcattcat gttattgata tttttcgtga tgatgggaat 900  
gtatataatag catatattgg tactgtattt aatagtacat ttaagaataa aaagcagttg 960  
gttattgttt atactatggg tgatgaacaa tcgcatgtgt atgattttat gcagattaca 1020  
gaagatatta gtgccatata tataaattct actgaaaata ttttagcaat aacgactatg 1080  
ggaagtgatt atcttgtaaa atatgagatt tcaaaattac agttaaattt agggattgtt 1140

gatcatgttg atgttataaa aattccacgt aatgtagtga aaaatattgc taatttttaca	1200
tatgtttgttg atacaatttt agggtttgat agtgttgaac atattaatat tcgtaatgta	1260
ttagctacaa aatcaactgt taattataag gtatctcagt tttttttaaa tatttagagaa	1320
atggaatttg gtgatttatt taagagttgg agtgggtgaat ataatgactt gttaataggt	1380
tatactatgc ctgctagtta tgggtgtaaata tatactacag aatattttaag cgatgttata	1440
actgtttcag gtaatgcagg ttttgtagag aagttcatat caactagtaa aatgggtgat	1500
gtattttaaga ttacagataa cttaattaat tatactagtg taaaccctac taatcatatg	1560
gcacatatga cattgcaatc aaaattgtca gatgggtgagg gtattacaga gcgtgcgggt	1620
aataaatcag ataattctgt aagtgaaggt ttagctacag gattgggttct tactagtaaa	1680
aatgatgatt tgtttaaaag tacagctagt cctattaatc atgcttttgg ttatgtaata	1740
aagcctactc gccatgtaac gcatgtaaca ttggaatcga agtcaccata tggtaaagag	1800
gttgtgaggc atatgaatcc taaaacggat aattctatac atacaagttc aataccaaga	1860
tcagtactga ctagtagaag cgatgatgta ttgaaaagta cagctagtcc tattaatcat	1920
gctttttggtt atgtaataaa gcctactcgc catgtaacgc atgtaacatt ggaatcgaag	1980
ttaccatattg ataaagaggt tgtgaggcat atgagtccta aaacggataa ttctatacat	2040
acaagttcaa taccaagatc agtactgact agtaaaagcg atgatgtatt gaaaagtaca	2100
gctagtccta ttaatcatgc ttttggttat gtaataaagc ctactcgcca tgtaacgcat	2160
gtaacattgg aatcgaagtt accatatgat aaagaggttg tgaggcatat gagtccataa	2220
acggataatt ctatacatc aagttcaata ccaagatcag tactgactag tagaagcgat	2280
gatgtattga aaagtacagc tagtcctatt aatcatgctt ttgggttatgt aataaagcct	2340
actcactatg taacgcatgt aacattggaa tcgaagtcac catatggtaa agaggttgtg	2400
aggcatatga atcctaaaac ggataattct atacatacaa gttcaatacc aagatcagta	2460
ctgactagta aaagcgatga tgtattgaaa agtacagcta gtcctattaa tgatgctttt	2520
ggttatatga aacctgctag ttctattgta gtatcattag gtgatactga tgtttcaag	2580
caagtgaaaa gtgttagtaa tgttccagta tatcttactc ctacagtaag atcagtatta	2640
gtaggtgatg cgtatcatgt atctggtagt gaaaaagata gtattggaca tgaacaagat	2700
ttgggtcatg gtgatgttag taccgatgtt gtattgaaac taatgagtga taatgtatca	2760
aacaatatta gtaggcatgt aaatgattct ttagctataa aacataagat attaggtaaa	2820
aaaataaagt ataataaag gcgtagtact gttagatctg ctgttaatat tcgcaataaa	2880
agtacagtga gtacaagata tacatctcat ggcatacaag aggctaataa tatgaatgtt	2940



```

acattgttta atcctacaca gcataatatt agtagttata atggtagttt attaaatagt 3000
aattctgctt ttaatactga aagttctgtt gattataaag tagtaattgc agtaatatct 3060
agtatactgc ttatcttttt attattaggt ggatttaaatt gtataaagtg gtatttagca 3120
aagttgaata gaagaaggat gtctaataat gaacagggat ttgtgatatt taatttggat 3180
agtattcaaa gtagtgtttc tgggtgtgcaa gtgacagaag gtaccacatc tcgaatagag 3240
agtctattct ag 3252

```

```

<210> 14
<211> 1836
<212> DNA
<213> Ehrlichia ruminantium

```

```

<400> 14
atgttattca aaccgggttc accggttgcc aagattacag aatctctgca taaatcagtt 60
atatatgagt tgaatagagt accagaaata catcttaata catgtcattg tataggagct 120
acaataggta caaaacttga tatctggatt gataataagt caggtcacgc gtgtactcca 180
gtaggaacat ctttatttct tatggaatgt attataccca ctgctgtaat aaatcatcca 240
cgtaatatat ctttacaaaa gttgacacaa gtattgtcta gtcgcttttc aagaacacaa 300
ccacttaagg ctgatgtata ttttatttga tcagaggaag aattcgagaa tttcagaagt 360
acagtatccc ctttatgtag tatgggactt aatgaacttt tacctgttta taatattggg 420
aaattcggag cattttgcgt atgtaggcca aaaagtatag gtaatagagg tgtagatgta 480
ctatttgatg aatacaaagc ttttaagggt ttaggaggtc tagaggattc taattttttt 540
aaaacaccct tatcaacctc taataccaca aaacgtaata caaaacaaag cacagcaaat 600
aatagagaac aaaaatttgt agtaactggg aagaaaattc aaagcaaaat acaaagtata 660
aaacatctac ataaaatatt ttctagatct tctactacac aatgttcacc tttaagtaca 720
ccagtcaata ctaaaacaca acataatata gaagaaaaaa cagcaagtag tacgcaagaa 780
ccaaatatcc aaaaggttat agtaactagc aatcaacctc atagagaaaa aacacaactt 840
atatgtacaa agtttcctga ggctccaaaa tatccatctt taaatcaaag acaagagaca 900
ggaggcaagt atttagaaca gcgtctatca aaaagtacag cagatagtag gccatttaca 960
caaaaaggta caacagattc tcaacaagtt gttagaccaa aaacacaatt tgcatctagt 1020
cctttttatt ttatcaaga acagccactt ttaactacaa aacattcatc ttcaaataca 1080
agacaagaga caggaggcaa gtatttagaa cagcgccctat caaaaagtac agcagatagt 1140
acgccattta cacaaaaagg tacaacagat tctcaacaag ttgttagacc aaaaacacaa 1200
tttgcaccta gtccttttta tttttatcaa gaacagccac ttttaactac aaaacattca 1260

```

tcttcaaadc aaagacaaga gacaggaggc aagtatttag aacagcgcct atcaaaaagt 1320  
 acagcagata gtacgccatt tacacaaaaa ggtacaacag attctcaaca agttgttaga 1380  
 ccaaaaacat ttgcatctag ttctttttat aaagaatcgg cacttgcaat tacaaagcat 1440  
 ccatcttcaa atcaaagaca agaggcaaca aacaagcatt tagaacagcg tccatcaaca 1500  
 aaaagtacag tagatcctca acaagttggt aggccaaaaa cacaatctgc atctagtcgt 1560  
 gtttataaag aacagggact tccaactaca aaacataaaa tattaagtgc tataaaagaa 1620  
 tctacagata gtagtacatc agttaatata ttaagttctg aagaagattt acggttttta 1680  
 aatgtagatt attctagtag ttgtgaaata ttatacgata ctttcagaga atcttataga 1740  
 gttagtgcct taccaacatc acctctcatc ccatcacatc aacttgaaga tgatgtattt 1800  
 gttgaagatg gttatcctcg tgcatcttat ctttga 1836

<210> 15  
 <211> 3570  
 <212> DNA  
 <213> Ehrlichia ruminantium

<400> 15  
 atgccactta cttttgatct atatgcatat gaaagaaaat taaattttct tctatgcaat 60  
 gctgtaaatt ctaatcctaa gttagttaat gtaataaatg tagtttgtgt aggatatact 120  
 gatgaaaata atcagttatt acttgctact gactacaaca ttccaccaga attacaccct 180  
 attcctagaa atcaatcctt atttcgtata aatgctaata taaaaactag cattataaca 240  
 aattctttta gattatctca agagtttgct ttaacacaag aggagctaaa taatggtaat 300  
 gttaaagtgtg aaatgtattg tttagtaggt aatgaaaatc ttgatgatit tactaaaata 360  
 tgtagtaagc ataaagtaag atataaaaat ctaacaacaa tttccaagaa tatttacaca 420  
 caattattta cagcagatat attaaaattt cctattaaca tagaacatgc tctatcaaat 480  
 atagcaaatt taaatgcaca atatatatat gcatctgac taataaatga atctgatcta 540  
 ataaatgcgt ctgatctaatt aaatgcgtct gatctaataa aagaagaaaa aattaagaat 600  
 attagaggaa gtactagtat attatatgat gcaatatgca gtacatatgc aactaatgat 660  
 taccatgtac tttctgtaaa atgcatagga tatactcata ataatcgaca actcatagtt 720  
 cacactcaat gtccagagaa ctttttacct atacctcaaa gtaactctct atttattgta 780  
 tgtgttgata tatcaccaga tatcataaca aataatgaaa atttatcctc tacttttgaa 840  
 ttaacagaaa gtgaaagtaa acaaagtact atcaattgtg caatgtactg tttagttaat 900  
 gatgaacaac ttggaagttt tactcataaa tgtaatacta caaataataa accaaagctt 960  
 caagatatta ttcaattttg ttctgtaata tgtataaac tcaatacaga aagaatatca 1020

tcattacaaa ttagcgaaga agagctaata aatagtgtag gaataggtga tgtaacattc	1080
agaaatttta gtgatctacg taaggaaaaa ctttaagaaaa tacagcaaata aaagaatgaa	1140
ctatgtagtgt caatatgcag tatatatgca gctaataact accatgtact ttctgtaaaa	1200
tgcataggat atactcataa taatcaacaa ctcatagttc aactcaatg tccagacagc	1260
cttttaccta tacctcaaag taactctcta tttattgtaa atgttgatgt atcaccagat	1320
atcataacaa ataataaaaa attatcctct acttttgcag taacagaaaag tgaaagtaag	1380
caaagtactc tcaagtgtgc aatgtactgt ttagttaatg atgaacaact tgaaagtttt	1440
actcataaat gtgatattac aaataataaa ccaaggcttc aagatattat tcaattttgt	1500
tctgtaatat gtataacact caatacagaa agaatgttat cattacaaat tagcgaagaa	1560
gagctaataa atagtgtagg aataggtgat gtaacattca aaaatttttag tgatctacgt	1620
caggaaaaat ttaataaaat acagcaaaca aataatgaac tatgtagtgc aatatgcatt	1680
tcacctgaag aaaataaaat aattgatata aaatgcgtag gacacactac cgctaagaat	1740
aaattagtag ttcatactga atgtccacta gctcttcttc ctacacctca aggtgattca	1800
ttattttcta tactgatggc tataccatac gctattatag caaataatgc catattatct	1860
cctgctttta aagtagtaaa aatgatctt ggtattaata gtaattatat tttatgcact	1920
gcatactgtc tagtaactaa gcatgatctt caagatttta ctaatgaagt gtcacggat	1980
ggtgcaatag gtgatagtat acaacaaaaa cgtcaaaaat ttgaaagtat cattaaatta	2040
tgttctgtaa aatgtgtaac attacatata caagaaatac tgtcattaaa tattagtcaa	2100
aaagaactaa tcaacgatat agggttatgt aatgcaactt tcaaataattt aagtaatctg	2160
catcaagaaa agattgatct acttaaacaa gtaaataata aattatgtcg tgaaatatgt	2220
aataaactta ggaaacataa aacacaatat ataagatgta taggaaatac tgttaatact	2280
aaattagtag ttaccactca gtgtccacga gatcttcttc cttttcctaa aggtcaatct	2340
ttattcatta taaggataaa tatatcacct aacattatat tacacagtaa aacactacgt	2400
aatacattta aattaacaac aagtgaaga tcagatcacc acattaaatg tgatatgtat	2460
tgcttagtat atgaagaaaa tattaagagt tttattgatg tatgtgatga tccaaataag	2520
ccatatattg aagagttaat tcaatattgt tctgtaaaat gtataaaatt gtatacacia	2580
gaaatgttat cattaaacat tagtgaagaa caactaatca acgatatagg gttatgtaat	2640
gcagaattta aatatgttga aagtaaacad ataattgaat cagtattgga cgcatttaatt	2700
tatatcgaaa tacaagcaaa taaactccta tgcggaattt tgcctacact ttgtgcttta	2760
tataaaaaag attttctcat caaaaagata cgttgatatag gtaatactat agatcctgaa	2820
caaggattaa caatttatcc ttctagtata taccocaaagg aattcttacc aactgcccaa	2880

```

gggtacatcctt tattttttaat acgaactagg atattaactg aagttatatt aagtactcct 2940
gaactagtga atgtacatat tctaaatgat gaagaaatgt tgaataagta tttattatgt 3000
gatatatatt gcctagtaga tgagaaaaac cttagaatat ttaagaatct ttgtacaaaa 3060
gcaagaaatc tttcagatat gataattaca tgtggtgtaa agtatgtag gatacataca 3120
aaagattcta aaagatttcc atttgatgaa gcaaaggat taaaacactt aggaggtata 3180
gacggaagat atctcgacga aggagatttt gacaaattac ttagttctgg actttatacc 3240
aaatcatcaa gtaagtcttc atcaacaata tcgactgaag aagaatcaag tacacaagaa 3300
gggaccata taaaacgtag tttaagatca acattattaa aaataagaaa acaaatagga 3360
cctgagtcctt catcatctgc tacattctca agtggagatg agttagattc agaagacgaa 3420
cttcaagaaa gaagacaaaa aagacgtgca agattagcaa gactacaaca tgaagaatca 3480
caaacaacaa aaagtaaaac aggaataggt ggtatcttgt ctgatcaaga agtttcacat 3540
cataaatctc aaagtaaaga tttagattag 3570

```

```

<210> 16
<211> 21
<212> DNA
<213> Ehrlichia ruminantium

```

```

<400> 16
atgagtcaca gttttattga g 21

```

```

<210> 17
<211> 21
<212> DNA
<213> Ehrlichia ruminantium

```

```

<400> 17
cactcaaaat cacaagaagt a 21

```

```

<210> 18
<211> 25
<212> DNA
<213> Ehrlichia ruminantium

```

```

<400> 18
atgtatttag tctatttagt agctg 25

```

```

<210> 19
<211> 23
<212> DNA
<213> Ehrlichia ruminantium

```

```

<400> 19
ataacatcta attgaacaat atc 23

```

<210> 20  
<211> 20  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 20  
atgaaaggat ctttatctgc 20

<210> 21  
<211> 20  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 21  
ccttcttctt cttcattatg 20

<210> 22  
<211> 19  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 22  
aagaattaca tgatgcagc 19

<210> 23  
<211> 22  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 23  
tcttctcttg ttatactctc tg 22

<210> 24  
<211> 23  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 24  
atggatttaa ataaactaat aaa 23

<210> 25  
<211> 19  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 25  
gcattttctc tacctacga 19

<210> 26  
<211> 26  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 26

gtacatagta tgtctttata taaaag 26

<210> 27  
<211> 23  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 27  
ccaaatatat aaatgatcta ttc 23

<210> 28  
<211> 25  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 28  
tccaccagag atgttatttg taaag 25

<210> 29  
<211> 25  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 29  
caacagaact ttcagtatta aaagc 25

<210> 30  
<211> 25  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 30  
gttaagtgtg aaatgtattg tttag 25

<210> 31  
<211> 25  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 31  
cactttctgt taattcaaaa gtaga 25

<210> 32  
<211> 25  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 32  
gtaggccaaa aagtataggt aatag 25

<210> 33  
<211> 25  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 33  
caacaaatac atcatcttca agttg

25

<210> 34  
<211> 25  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 34  
agggttactt attgtagtca gagtg

25

<210> 35  
<211> 25  
<212> DNA  
<213> Ehrlichia ruminantium

<400> 35  
cctcttcgta tacaggatta ccatt

25

## INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2006/003870

## A. CLASSIFICATION OF SUBJECT MATTER

INV. C12N1/20 C12N15/09 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, EMBASE, BIOSIS, WPI Data, PAJ

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2006/045338 A (AGRONOMIQUE POUR LE DEV CIRAD [FR]; CENTRE NAT RECH SCIENT [FR]; FRUTO) 4 May 2006 (2006-05-04) cited in the application page 2, line 19 - page 14, line 28; sequences 1-6, 9, 11, 13, 20	1-3
A	FRUTOS R ET AL: "Comparative genomic analysis of three strains of Ehrlichia ruminantium reveals an active process of genome size plasticity" JOURNAL OF BACTERIOLOGY 2006 UNITED STATES, vol. 188, no. 7, April 2006 (2006-04), pages 2533-2542, XP002430891 ISSN: 0021-9193 cited in the application the whole document	1-3



Further documents are listed in the continuation of Box C.



See patent family annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

24 April 2007

Date of mailing of the international search report

09/05/2007

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Donath, Cornelia



## INTERNATIONAL SEARCH REPORT

International application No

PCT/IB2006/003870

## C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	BEKKER CORNELIS P J ET AL: "Transcription analysis of the major antigenic protein 1 multigene family of three in vitro-cultured Ehrlichia ruminantium isolates" JOURNAL OF BACTERIOLOGY, vol. 187, no. 14, July 2005 (2005-07), pages 4782-4791, XP002430892 ISSN: 0021-9193 the whole document	1-3
A	COLLINS NICOLA E ET AL: "The genome of the heartwater agent Ehrlichia ruminantium contains multiple tandem repeats of actively variable copy number" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, vol. 102, no. 3, 18 January 2005 (2005-01-18), pages 838-843, XP002430893 ISSN: 0027-8424 the whole document	1-3
A	ALLSOPP M T ET AL: "Ehrlichia ruminantium major antigenic protein gene (map1) variants are not geographically constrained and show no evidence of having evolved under positive selection pressure" JOURNAL OF CLINICAL MICROBIOLOGY, WASHINGTON, DC, US, vol. 39, no. 11, November 2001 (2001-11), pages 4200-4203, XP002321870 ISSN: 0095-1137 the whole document	1-3

## INTERNATIONAL SEARCH REPORT

### Information on patent family members

International application No

PCT/IB2006/003870

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2006045338	A	04-05-2006	NONE